

Relations between Electromagnetic Form Factors of Baryons

A. J. Buchmann^{a*†}

^aInstitute for Theoretical Physics, University of Tübingen, Auf der Morgenstelle 14
D-72076 Tübingen, Germany

The inclusion of two-body exchange currents in the constituent quark model leads to new relations between the electromagnetic properties of octet and decuplet baryons. In particular, the $N \rightarrow \Delta$ quadrupole transition form factor can be expressed in terms of the neutron charge form factor.

1. Neutron and Δ charge form factors

In Ref. [1] we have shown that the Sachs charge form factor $G_E^n(q^2)$ and charge radius $r_n^2 = -6(d/dq^2)G_E^n(q^2)|_{q^2=0}$ of the neutron are dominated by quark-antiquark pair exchange currents shown in Fig.1(b-c). The latter describe the gluon and pion degrees of freedom, while the one-body currents in Fig.1(a) describe the valence quark degrees of freedom in the nucleon. The two-body exchange charge operator contains a spin-dependence of the schematic form $\rho_{[2]} \propto \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j Y^0(\mathbf{q}) - \frac{\sqrt{6}}{2} [(\boldsymbol{\sigma}_i \times \boldsymbol{\sigma}_j)^2 \times Y^2(\mathbf{q})]^0$. This gives different matrix elements for quark pairs in spin 0 and spin 1 states. Because a down-down pair in the neutron is always in a spin 1 state, while an up-down pair can be in spin 0 or spin 1 states, an asymmetry in the charge distribution of up and down quarks arises. Consequently, a nonvanishing neutron charge form factor (see Fig.2) and radius is obtained [1]:

$$r_n^2 = -\frac{b^2}{3m_q} \left(\delta_g(b) + \delta_\pi(b) \right) = -b^2 \frac{M_\Delta - M_N}{M_N}. \quad (1)$$

Here, b is the quark core (matter) radius of the nucleon, m_q is the constituent quark mass, and the functions $\delta_g(b)$ and $\delta_\pi(b)$ are the gluon and pion contributions to the N - Δ mass splitting, which satisfy $M_\Delta - M_N = \delta_g(b) + \delta_\pi(b)$.

The spin-spin term in the charge operator is also responsible for the following relation between the charge form factors of the nucleon and Δ :

$$G_E^p(q^2) - G_E^{\Delta^+}(q^2) = G_E^n(q^2), \quad r_p^2 - r_{\Delta^+}^2 = r_n^2, \quad (2)$$

where $r_{\Delta^+}^2$ is the charge radius of the Δ^+ and $\mathbf{q}$ is the three-momentum transfer of the photon. An analogous result holds for the difference of neutron and Δ^0 form factors. The charge form factor of the Δ^0 and the corresponding charge radius are zero in the present

*email:alfons.buchmann@uni-tuebingen.de

[†]Talk given at Intern. Symposium on Physics of Hadrons and Nuclei, Tokyo, Japan, 14-17 Dec., 1998, published in Nucl. Phys. **A670** (2000) 174c-177c.




Figure 1. Feynman diagrams of the four vector current $J^\mu = (\rho, \mathbf{J})$: photon (γ) coupling to (a) one-body current $J_{[1]}^\mu$, and to (b-d) two-body gluon and pion exchange currents $J_{[2]}^\mu$. Diagrams (b-d) must be taken into account in order to satisfy the continuity equation $q_\mu J^\mu = 0$ for the electromagnetic current J_μ . They represent the nonvalence (gluon and pion) degrees of freedom in the nucleon.

model as it should be on general grounds [3]. From its derivation [4] it is evident that the charge radius relation is independent of whether gluon or pion exchange is the dominating residual interaction between constituent quarks. Using a general parametrization method, Dillon and Morpurgo [5] have recently shown that, if three-quark currents and strange quark loops are neglected, $r_p^2 - r_{\Delta^+}^2 = r_n^2$ is a consequence of the symmetries and dynamics of QCD that is model-independently valid. They have also shown that three-body currents slightly modify, but do not invalidate the general relation between the proton, neutron, and Δ charge radii.

We repeat that Eq.(2) is a consequence of the spin-spin $\boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j / (m_i m_j)$ term in $\rho_{[2]}$, which leads to a Δ charge radius that is larger than the proton charge radius. This effect is of the same generality as, and closely connected with the $N - \Delta$ mass splitting due to the spin-spin interaction in the Hamiltonian. The latter is repulsive in quark pairs with spin 1 and makes the Δ heavier than the nucleon.³

2. Electromagnetic $N \rightarrow \Delta$ transition form factors

In the constituent quark model with exchange currents a connection between the neutron charge form factor $G_E^n(\mathbf{q}^2)$ [1] and the $N \rightarrow \Delta$ quadrupole transition form factor $F_Q^{p \rightarrow \Delta^+}(\mathbf{q}^2)$ [2,4] emerges:

$$F_Q^{p \rightarrow \Delta^+}(\mathbf{q}^2) = -\frac{3\sqrt{2}}{\mathbf{q}^2} G_E^n(\mathbf{q}^2), \quad Q_{p \rightarrow \Delta} = \frac{r_n^2}{\sqrt{2}}, \quad r_{Q, p \rightarrow \Delta^+}^2 = \frac{11}{20} b^2 + r_{\gamma q}^2. \quad (3)$$

The $N \rightarrow \Delta$ quadrupole transition form factor is a measure of the intrinsic deformation of the nucleon and the Δ . The above results for the $N \rightarrow \Delta$ transition quadrupole moment, $Q_{p \rightarrow \Delta^+}$, and the transition quadrupole radius⁴, $r_{Q, p \rightarrow \Delta^+}^2$, were derived before[2,4]. They are seen here to be the 0th and 1st moment of the more general relation between G_E^n and

³Combining Eq.(1) and Eq.(2) yields $r_{\Delta^+}^2 - r_p^2 = [b^2/(3m_q)](M_{\Delta^+} - M_p)$, i.e., a relation between the mass difference between Δ^+ and proton and a corresponding charge radius difference.

⁴The term $\frac{1}{4}b^2$ in Eq.(53) of Ref.[2] should be replaced by $\frac{11}{20}b^2$




Figure 2. Neutron charge form factor $G_E^n(Q^2)$ and the quadrupole transition form factor $F_Q^{p \rightarrow \Delta^+}(Q^2)$ as a function of four-momentum transfer $Q^2 = -q^2$. The crosses, triangles, circles are the recent data [6]. The upper curve is a quark model calculation with exchange currents [1,2]. The gluon and pion contributions to $G_E^n(Q^2)$ are shown separately. The lower curve is $F_Q^{p \rightarrow \Delta^+}(Q^2) = -3\sqrt{2} G_E^n(Q^2)/Q^2$.

$F_Q^{p \rightarrow \Delta^+}$, plotted as the lower curve in Fig.2. The quark model with exchange currents explains $Q_{p \rightarrow \Delta^+}$ as a *double spin flip* of two quarks, with all valence quarks remaining in the dominant, spherically symmetric $L = 0$ state. The double spin-flip comes from the tensor term in $\rho_{[2]}$. The latter is closely related to the tensor term in the Hamiltonian. The quark core (D waves in the nucleon) also contributes to $Q_{p \rightarrow \Delta^+}$. This valence quark contribution amounts to about 20% (due to the smallness of the D wave amplitudes) of the double spin flip amplitude [4]. We conclude that the collective gluon and pion degrees of freedom are mainly responsible for the deformation of the N and Δ .

Due to the first relation in Eq.(3), the quadrupole transition radius can also be expressed as the 2nd moment of $G_E^n(\mathbf{q}^2)$, namely, $r_{Q,p \rightarrow \Delta^+}^2 = (18/r_n^2) (d/d\mathbf{q}^2)^2 G_E^n(\mathbf{q}^2)|_{\mathbf{q}^2=0}$. Because the quark core radius b is fixed by Eq.(1), one could extract the charge radius of the light constituent quarks, $r_{\gamma q}^2$, from both the $G_E^n(\mathbf{q}^2)$ data, and from the slope of $F_Q^{p \rightarrow \Delta^+}(\mathbf{q}^2)$ at $\mathbf{q}^2 = 0$. Both determinations of $r_{\gamma q}^2$ should agree.

It is interesting that the additive quark model relation between the magnetic $N \rightarrow \Delta$ transition and the neutron magnetic moments $\mu_{p \rightarrow \Delta^+} = -\sqrt{2}\mu_n$ remains unchanged after including the gauge-invariant two-body exchange currents of Fig.1(b-d); and that it continues to hold even at finite momentum transfers

$$F_M^{p \rightarrow \Delta^+}(\mathbf{q}^2) = -\sqrt{2} G_M^n(\mathbf{q}^2), \quad \mu_{p \rightarrow \Delta^+} = -\sqrt{2}\mu_n, \quad r_{M,p \rightarrow \Delta^+}^2 = -\sqrt{2} \frac{\mu_n}{\mu_{p \rightarrow \Delta^+}} r_{M,n}^2, \quad (4)$$

where $r_{M,p \rightarrow \Delta^+}^2$ is the magnetic $N \rightarrow \Delta$ transition radius, and $r_{M,n}^2$ the magnetic radius of the neutron. The transition magnetic moment predicted by Eq.(4) underestimates the empirical value by 30%. This discrepancy between theory and experiment can presumably be explained by including spatial three-body currents in the theoretical description [7].

Combining Eq.(3) and Eq.(4) we find that the ratio of the charge quadrupole and magnetic dipole $N \rightarrow \Delta$ transition form factors can be expressed in terms of the experimentally better known elastic neutron form factors

$$\frac{F_Q^{p \rightarrow \Delta^+}(\mathbf{q}^2)}{F_M^{p \rightarrow \Delta^+}(\mathbf{q}^2)} = \frac{3}{\mathbf{q}^2} \frac{G_E^n(\mathbf{q}^2)}{G_M^n(\mathbf{q}^2)}, \quad \frac{C2}{M1} = \frac{M_N \omega_{cm}}{2 \mathbf{q}^2} \frac{G_E^n(\mathbf{q}^2)}{G_M^n(\mathbf{q}^2)}, \quad (5)$$

where $\omega_{cm} = 258$ MeV is the center of mass energy of the photon-nucleon system at the Δ resonance. For example, this yields $\frac{C2}{M1}(\mathbf{q}^2=0) = -1.04 r_n^2/(2 \mu_n) = -0.030$ and $\frac{C2}{M1}(\mathbf{q}^2=4.2 \text{ fm}^{-2}) = -0.042$. Sign and magnitude of these theoretical predictions are in agreement with recent experimental data $\frac{C2}{M1}(\mathbf{q}^2=4.2 \text{ fm}^{-2})_{exp} = -0.046(8)$ [8].

3. Relations between octet and decuplet hyperon charge radii

Using the general parametrization method of Refs.[5,7], we find the following relations between octet and decuplet charge radii for strange hyperons:

$$r_{\Sigma^-}^2 - r_{\Sigma^{*-}}^2 = r_{\Xi^-}^2 - r_{\Xi^{*-}}^2 = r_n^2 (x + x^2), \quad r_{\Sigma^+}^2 - r_{\Sigma^{*+}}^2 = r_{\Xi^0}^2 - r_{\Xi^{*0}}^2 = r_n^2 (2x - x^2), \quad (6)$$

where $x = m_u/m_s$ is the ratio of nonstrange to strange quark masses. Again, it is the $\boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j/(m_i m_j)$ term in the charge operator that leads to Eq.(6).

In summary, by including two-body currents in the constituent quark model we have found hitherto unknown relations between the electromagnetic form factors of octet and decuplet baryons. In particular, the $C2/M1$ ratio in the electromagnetic $N \rightarrow \Delta$ transition can be expressed in terms of the elastic form factors of the neutron.

REFERENCES

1. A. Buchmann, E. Hernández, and K. Yazaki, Phys. Lett. **B269**, 35 (1991); Nucl. Phys. **A569** (1994) 661.
2. A. J. Buchmann, Z. Naturforsch. **52a**, 877 (1997).
3. S. Coleman, Aspects of Symmetry, Cambridge University Press, 1985.
4. A. J. Buchmann, E. Hernández, and A. Faessler, Phys. Rev. **C55** (1997) 448.
5. G. Dillon and G. Morpurgo, Phys. Lett. **B448** (1999) 107.
6. C. Herberg *et al.*, Eur. Phys. J. **A5** (1999) 131; J. Becker *et al.*, *ibid.*; M. Ostrick *et al.*, Phys. Rev. Lett. **83** (1999) 276; I. Passchier *et al.*, Phys. Rev. Lett. **82** (1999) 4988; for refs. to the other data see Ref.[2].
7. G. Morpurgo, Phys. Rev. **D40** (1989) 2997.
8. P. Bartsch *et al.*, Proceedings of Baryons' 98, Bonn, Germany, World Scientific, 1999, eds. D. W. Menze and B. Metsch, pg. 757; M. O. Distler, *ibid.*, pg. 753; R. Gothe, *ibid.* pg. 394.